

## Effectiveness of Using Toothpaste and Mouthwash of Beluntas Leaf Ethanol Extract in Reducing *Streptococcus mutans* Bacteria Number in Tooth Plaque

Maria Martina Nahak<sup>1\*</sup>, Regina Tedjasulaksana<sup>1</sup>, I Gusti Ayu Raiyanti<sup>1</sup>

1. Department of Dental Hygienist, Polytechnic of Health-Ministry of Health, Denpasar, Indonesia.

### Abstract

The aim of this study is to analyze the differences in the effectiveness of toothbrushing and gargling using 10% beluntas leaf ethanol extract toothpaste and mouthwash to reduce the number of colonies of *Streptococcus mutans* in dental plaque.

The type of this research is experiment with completely randomized with pre-posttest control group design, with 105 samples divided into three treatment groups and four control groups. There is a significant decrease in the amount of *Streptococcus mutans* in dental plaque after receiving treatment at 95% confidence level with  $P=0.00$  or  $P<0.05$  and there was no difference in effectiveness of toothbrushing and gargling using 10% ethanol extract of leaves beluntas with 0.12% chlorhexidine and 1% triclosan to decrease the amount of *Streptococcus mutans* bacteria in dental plaque, with  $P= 0.166$  or  $P> 0.05$ .

Both beluntas leaf ethanol extract gargle and toothpaste group with positive control group using chlorhexidine and triclosan, is equally effective in reducing the number of colonies of *Streptococcus mutans* in dental plaque, so it can be concluded that the extract of beluntas leaves ethanol effectively decreases the amount of *Streptococcus mutans* colonies in dental plaque and there is no difference in effectiveness between the treatment group and the control group.

Experimental article (J Int Dent Med Res 2018; 11(3): 830-833)

**Keywords:** 10% beluntas leaf ethanol extract toothpaste, 10% beluntas leaf ethanol extract gargle, Tooth plaque, *Streptococcus mutans*.

**Received date:** 26 March 2018

**Accept date:** 21 May 2018

### Introduction

Caries is a disease of tooth hard tissue, most commonly found in the oral cavity, can affect all populations regardless of age, gender, race or socio-economic circumstances and is a leading cause of tooth loss.<sup>1</sup> Caries is caused by microorganism activity by distributing carbohydrates in the mouth, characterized by demineralization of inorganic materials followed by organic material damage from enamel and dentin. Demineralization of hard tissue occurs in the crown of the email and dentin and the root part of cementum and dentin.<sup>2</sup>

Caries should be addressed by curative effort that is by treating and patching all the teeth

that hollow. Promotional efforts are done by providing correct information and education about how to maintain dental health.<sup>3</sup> Mechanical preventive measures are performed by brushing, while chemical methods are performed by gargling with antiseptics such as chlorhexidine, or mouthwash from plant extracts to reduce plaque accumulation. 0.12% chlorhexidine, a broad-spectrum antimicrobial, is more effective against Gram-positive bacteria,<sup>4</sup> but may have adverse side effects such as staining on teeth and restorations although not permanent.<sup>5</sup> Prevention of plaque accumulation using plant extracts or traditional medicines has been widely studied today, one of which is the *beluntas* leaf which can inhibit the growth of *Streptococcus mutans* invitro.<sup>6</sup> *Beluntas* leaves contain active compounds such as alkaloids, flavonoids, triterpenoids, tannins, and phenols as well as other essential oil derivatives,<sup>7,8</sup> which work synergistically to inhibit the growth of *Streptococcus mutans*.

Mouthwash is a solution or liquid used to rinse the oral cavity with a number of purposes: to exclude pathogenic bacteria, work as an

#### \*Corresponding author:

Maria Martina Nahak  
Department of Dental Hygienist,  
Polytechnic of Health-Ministry of Health,  
Denpasar, Indonesia.  
E-mail: marianahak@rocketmail.com

astringent, to remove odor, remove infection and prevent caries, as a topical anti-inflammatory and analgesic agent.<sup>9</sup> The mouthwash composition consists of three main components: the active ingredient, the solvent, and the surfactant which serves to remove debris from the tooth and dissolve the other material.<sup>9</sup> The mouthwash is effective for reaching the most difficult place to clean with a toothbrush but its use cannot be as a toothbrush substitution.<sup>9</sup>

Mouthwashes as well as toothpastes have functions can be categorized as cosmetic and or therapeutic.<sup>9</sup> Toothpaste is a dental cleaning agent containing abrasive, detergent and one or more therapeutic substances,<sup>10</sup> which are useful for maintaining oral health.

According to Soho Global Health's Sales and Marketing for Professional Products Vice President, In line with the back to nature trend, herbal products have been used by 80 percent of the world's population, while in Indonesia it always increases year by year and in 2012 reaches 13 trillion rupiah or about 2% of the total market of herbal medicine in the world and this makes drug manufacturers interested in continuing to develop herbal medicine.<sup>11</sup> Beluntas leaf ethanol extract also can be used as one of the active ingredients in mouthwash and toothpaste as a therapeutic agent that serves to prevent plaque accumulation and in the future can be developed into toothpaste products and herbal mouthwash that can be marketed in Indonesia.

### Materials and methods

This research is experiment with completely randomized with pre-posttest control group design. The sample size is 105 respondents, consisting of 3 treatment groups and 4 control groups. How to select the sample is simple random sampling. The research procedure is as follows: a). At first swab plaque using sterile copan swab on the surface of the buccocervical of the maxillary tooth. Results of plaque swabs were incubated for 24 hours at 37 °C, then diluted according to preliminary test results, reproduced in Mueller Hinton solid media on blood agar plates, grounded for 24 hours at 37 °C; b). Conducted isolation to separate the bacteria *Streptococcus mutans* from other bacterial colonies; c). Identification test with Gram test, catalase test, oxidase test and

identification test using Streptococcus grouping kit to confirm the presence of *Streptococcus mutans* bacteria and calculate the number of *Streptococcus mutans* manually. The calculated result is multiplied by the plaque dilution; d). Respondents in the treatment group: brushing or gargling with 10% *beluntas* leaf ethanol extract tooth paste and mouth wash or brushing and gargling with 10% *beluntas* leaf ethanol extract tooth paste and mouthwash for 7 days consecutively; e). Respondents in the negative control group, brushing or gargling with a toothpaste base, or a mouthwash base, while for the positive control group, toothbrushes use 1% triclosan toothpaste and gargle with 0.12% chlorhexidine, for 7 days consecutively; f). On the eighth day, swabs were performed on all respondents, all swabs were treated as pretest stage and then counted the number of *Streptococcus mutans* bacteria manually. The calculated result is multiplied by dilution.

Collected data were analyzed statistically by SPSS, using descriptive test, normality test using Shapiro-Wilk test and homogeneity test using Levene test. Normality and homogeneity test results showed normal and non-homogeneous distribution data, comparative test between groups of data was done by Repeated Measure test followed by Least Significance Difference test to see which groups were significantly different.

### Results

Sample group	Number of <i>Streptococcus mutans</i> before Treatment (Pretest) (CFU/ml)				Number of <i>Streptococcus mutans</i> after Treatment (Posttest) (CFU/ml)			
	Min	Max	Mean	SD	Min	Max	Mean	SD
P <sub>0</sub>	9	293	141.8	93.30	4	300	143.1	89.74
P <sub>1</sub>	8	300	102.3	103.80	1	300	95.7	90.40
P <sub>2</sub>	9	305	67	72.07	0	201	41.9	70.49
P <sub>3</sub>	19	325	150.1	105.03	0	305	43.7	79.03
P <sub>4</sub>	2	211	66.7	66.55	1	114	26.6	30.89
P <sub>5</sub>	13	300	102.3	105.18	0	81	31.5	29.91
P <sub>6</sub>	6	305	146.3	120.23	0	300	103.3	96.48

**Table 1.** The Effectiveness of Tooth Brushing and Gargling Using *beluntas* leaf ethanol extract Tooth Paste and Mouthwash to Reduce Number of *Streptococcus mutans* Bacteria in Dental Plaque.

Table 1 above shows that almost all sample group, except P<sub>0</sub>, a decrease in the number of *Streptococcus mutans* bacteria after treatment.

The results of the Repeated Measure test are shown in table 2 below:

Effect		Value	F	p
Pre and post test	Pillai's Trace	0.121	13.503	0.000
	Wilks' Lambda	0.879	13.503	0.000
	Hotelling's Trace	0.138	13.503	0.000
	Roy's Largest Root	0.138	13.503	0.000
Effectiveness comparison of pretest and posttest material	Pillai's Trace	0.087	1.565	0.166
	Wilks' Lambda	0.913	1.565	0.166
	Hotelling's Trace	0.096	1.565	0.166
	Roy's Largest Root	0.096	1.565	0.166

**Table 2.** Effectiveness difference of Tooth Brushing and Gargling using *beluntas* leaf ethanol extract tooth paste and mouth wash to Reduce Number of *Streptococcus mutans* Bacteria in Dental Plaque.

Table 2 shows that the calculated F value on each test is the same, and selected variance test is Pillai's Trace because the data is not homogeneous. Visible value of Pillai's Trace = 0.121, with value of calculated F = 13.503 and p-value = 0.000 or  $P < 0.05$ , hence it is statistically proven that quantity of *Streptococcus mutans* bacteria before and after treatment in all admission and control group, significantly different and significant with 95% confidence interval.

The comparison of effectiveness between the test materials used, the value of Pillai's Trace = 0.087, has the value of calculated F = 1.565 and the value of  $P = 0.166$  ( $P > 0.05$ ), so there is no significant difference between test materials in decreasing the number of *Streptococcus mutans* in dental plaque.

## Discussion

Descriptive test results showed that there was a decrease in the number of *Streptococcus mutans* in all groups after treatment, except in group P<sub>0</sub>. The multivariate analysis showed that there was a significant difference between the number of bacteria before and after treatment. Multivariate analysis also showed that all types of mouthwashes or toothpaste used by the study subjects had a significant effect on reducing the amount of *Streptococcus mutans* bacteria and there is no type of mouth wash or toothpaste that was superior to others. These results indicate that all test materials have the same effect to decrease the amount of *Streptococcus mutans* in dental plaque. The results of this study are in accordance with the research conducted by

Nahak, et al. (2015), which found that gargling with 10%, 20% and 30% *beluntas* ethanol extracts and 0.12% chlorhexidine, could decrease the number of *Streptococcus* sp. on dental plaque.<sup>12</sup>

0.12% chlorhexidine mouthwash is an antiseptic that has antibacterial activity. Its ability to reduce aerobic or anaerobic bacteria in the oral cavity reaches 54-97%.<sup>3</sup> Mechanism of action is allegedly by damaging the bacterial cell wall, inhibits bacterial enzymatic system, removing bacterial lipopolysaccharide causing bacterial cell death.<sup>13,14</sup> 1% Triclosan toothpaste has the effect of significantly lowering the number of *Streptococcus mutans* in dental plaque. Triclosan is a broad-spectrum non-ionic antiseptic, used in a variety of products, such as soap, mouthwash and toothpaste. The antimicrobial activity was obtained at concentrations of 0.2-2% and was bacteriostatic. Triclosan works by affecting the structure and function of cytoplasmic membranes that cause cells lysis.<sup>15</sup>

The results of this study also showed that the formula of 10% *beluntas* leaf ethanol extract toothpaste and mouthwash could significantly reduce the amount of *Streptococcus mutans* bacteria in dental plaque. This is allegedly related to the content of bioactive substances. Tannin is a phenolic compound, has antibacterial activity by forming a complex with proline that is a kind of protein in bacterial cell wall, causing protein leakage, damaging bacterial cell wall causing bacterial cell death.<sup>16,17</sup> Steroids have antibacterial activity allegedly by causing leakage in the liquid part of *phosphatidyl ethanolamine* in liposome-rich bacteria.<sup>18,19</sup> Flavonoid compounds are a group of phenol compounds, have antibacterial activity by causing protein leakage, resulting in leakage of bacterial cell wall, inhibiting bacterial protein synthesis and possibly interfere with bacterial cell DNA function.<sup>18,20</sup> The actual working mechanism of the active substances contained in the *beluntas* leaf extract is not known with certainty, but it is suspected that the active substances are working synergistically to inhibit bacterial growth in plaque thus decreasing the amount of colony of *Streptococcus mutans* in dental plaque. Multivariate test results also showed that no test material had superior efficacy compared to the others to decrease the number of colonies of *Streptococcus mutans* in dental plaque.

In line with the “back to nature“ trend, the use of herbal medicine in the world continues to increase as well as in Indonesia.<sup>11</sup> Beluntas leaf ethanol extract containing antibacterial ingredients can be used as an active ingredient in mouthwash and toothpaste as a therapeutic agent that serves to prevent plaque accumulation and in the future can be developed into toothpaste products and herbal mouthwash that can be marketed in Indonesia.

## Conclusions

*Beluntas* leaf ethanol extract tooth paste and mouthwash with 10% concentration effective to significantly reduce the amount of *Streptococcus mutans* in dental plaque and no effectiveness difference between 10% *beluntas* leaf ethanol extract tooth paste and mouthwash compared with positive control in lowering the number *Streptococcus mutans* colonies in dental plaque or in other words, *beluntas* leaf ethanol extract tooth paste and mouthwash have the ability to decrease the amount of *Streptococcus mutans* colony in dental plaque comparable to 1% triclosan toothpaste and 0.12% chlorhexidine mouthwash.

## Acknowledgements

Obtained from the University Committee and Respondent agreement. Source of Funding: Indonesian Ministry of Health

## Declaration of Interest

The authors report no conflict of interest and the article is not funded or supported by any research grant.

## References

1. Darwita RR, Gityawati C, Andreas P, Setiawati F, Soekanto SA. The role of education for 21 Days in improving the effectiveness of brushing teeth of children age 5 – 8-year-old in Pembeliangan village North Kalimantan. *J Int Dent Med Res.* 2017;10:683-8.
2. Badruddini A, Kiptiyah NM, Prihartono N, Agtini MD, Musadad DA. The Association between Sweet Food Consumption, Time of Tooth Brushing and Dental Caries Experience in 12-to15 Year-old Children in Indonesia (Analysis of Indonesian Health Basic Research Data, 2013). *J Int Dent Med Res.* 2017;10:583-9.
3. Suahrsini M, Budiarto SB, Indarti IS, Rudianto YE, Widyagarini A. Effect of Tooth Brushing, Using Song and Dental Model, on Plaque Index of Children with Visually Impairment. *J Int Dent Med Res.* 2017;10(special issue): 608-11.
4. Shahani MN, Reddy VVS. Comparison of Antimicrobial Substantivity of Root Canal Irrigants in Instrumented Root Canals up to 72 Hours: An *In Vitro* Study. *Journal of Indian Soc. Pedod. Prev. Dent.* 2011;29: 28-33.
5. Peterson D. Family Gentle Dental Care. Article. 2011; (Serial Online). (cited 2011, Agustus 8). Available from: <http://www.dentalgentlecare.com/periguard.htm>
6. Nahak MM. Ekstrak Etanol Daun Beluntas (*Pluchea indica*. L.) dapat Menghambat Pertumbuhan Bakteri *Streptococcus mutans*. Laporan Tesis. 2012; Program Pasca Sarjana Universitas Udayana Denpasar
7. Dalimartha S. Atlas Tumbuhan Obat Indonesia. Jilid I. Trubus Agriwidya Jakarta. 1999:18-21
8. Nahak MM, Tedjasulaksana R, Dharmawati IGGA. Khasiat Ekstrak Daun Beluntas untuk Menurunkan Jumlah Bakteri pada Saliva. *Interdental Jurnal Kedokteran Gigi Denpasar.* 2007;5(3):139-42.
9. Susanto IGOA. Optimasi Obat Kumur (Mouthwash) Minyak Atsiri Sereh Dapur (*Cymbopogon citratus*). Skripsi. 2012. Surakarta: Fakultas Farmasi Universitas Muhammadiyah.
10. Midda M, Coosey MW. Clinical Uses of an Enzyme Containing Dentrifices. *Journal of Clinical Periodontal.* 1986; 13:950
11. Hardianti AL, Rini CL. Tren Penggunaan Obat Herbal Meningkatkan. Artikel. 2014.(online). (Cited 2 Agustus 2016). Available from: [http://www.republika.co.id/berita/gaya\\_hidup/info-sehat/](http://www.republika.co.id/berita/gaya_hidup/info-sehat/)
12. Nahak MM, Tedjasulaksana R, Sumerti NN. Efektivitas Kumur-kumur Ekstrak Etanol Daun Beluntas (*Pluchea indica*. L.) untuk Menurunkan Jumlah *Streptococcus* sp. pada Plak Gigi. *Jurnal Skala Husada* 2015; 12(1):56-64
13. Kuyyakanond T, Quenel LB. The Mechanism of Action of Chlorhexidine. *FEMS Microbiol Lett.* 1992;79(1-3): 211-215
14. Mandel ID. Antimicrobial Mouth Rinses: Overview & Update. *J Am Dent Assoc.* 1994;125(25): 2S -10S.
15. Loho T, Utami L. Uji Efektivitas Antiseptik Triclosan 1% terhadap *Staphylococcus aureus*, *Escherichia coli*, *Enterococcus faecalis* dan *Pseudomonas aeruginosa*. *Majalah Kedokteran Indonesia* 2007;57(6):171-8.
16. Machado TB, Pinto AV, Pinto MCFR, Leal ICR, Silva MG, Amaral ACF, Kuster RM, Netto-dos Santos KR. *In vitro* Activity of Brazilian medicinal plants, naturally occurring naphthoquinones and their analogues against Methicilline Resistant *Staphylococcus aureus*. *Int J Antimicrobial Agents* 2003;21:279-84.
17. Braga LC, Shupp JW, Cummings C, Jett M, Takahashi JA, Carmo LS, Chartone-Souza E, Nasclmento AMA. Pomegranate Extract Inhibits *Staphylococcus aureus* Growth and Subsequent Enterotoxin Production. *J Ethnopharmacol.* 2005;96 : 335-9.
18. Mohamed SSH, Hansi PD, Thirumurugan K. Antimicrobial Activity and Phytochemical Analysis of Selected Indian Folk Medicinal Plants. *Int J Pharma Sci Res.* 2010; 1(10): 430-4.
19. Epand RF, Savage PB, Epand RM. Bacterial Lipid Composition and the Antimicrobial Efficacy of Cationic Steroid Compounds (Ceragenins). *Biochimica et Biophysica Acta.* 2007; 1768(10): 2500-9
20. Hussain A, Wahab S, Zarin I, Sarfaraj Hussain M.D. Antibacterial Activity of the Leaves of *Cocconiaindica* (W. and A) Wof India. *Advances in Biological Research.* 2010;4(5): 241-8.